

AD-P008 481



Multivalent ligands for diagnosis and therapeutics

Shubb D. Sharma^a, Victor J. Hruby^a, Mac E. Hadley^b, Michael E. Granberry^c
and Stanley P.L. Leong^c

*Departments of ^aChemistry, ^bAnatomy and ^cSurgery, University of Arizona,
Tucson, AZ 85721, U.S.A.*

Introduction

Multivalent ligands have been shown to exhibit extraordinary properties such as: a) enhanced affinity towards interacting cellular binding sites, b) increased potency, c) stability towards proteolytic enzymes, and d) longer duration of activity [1,2]. Such properties result from simultaneous multiple interactions (cooperative affinity) between the ligand and the acceptor. We are developing a new class of multivalent peptide hormone (neurotransmitter) - macromolecular composites which may serve as powerful diagnostic, imaging, and therapeutic tools. Composites have been synthesized in which multiple copies of a biospecific ligand were covalently attached to a biologically compatible but inert polyfunctional macromolecule. In addition, the conjugation of multiple copies of a fluorophore directly to the macromolecule provided an enhanced visual means of detection of ligand-macromolecular composites bound to target cells in *in vitro* binding assays. A fluorescent melanotropin (MSH) - macromolecular composite has been synthesized and used to demonstrate the presence of specific melanotropin receptors on various human melanoma cell lines.

Results and Discussion

The composition of various composites synthesized in this study are described in Table 1. Poly-lysine or Poly-vinyl alcohol used as a substrate was derivatized either with 4-(*p*-maleimidophenyl)butyric acid-*N*-hydroxysuccinimide ester, or with 3-(2-pyridyldithio)-propionic acid-*N*-hydroxysuccinimide ester for the synthesis of composites in which the MSH molecules are attached to the polymer via a thioether or a disulfide linkage, respectively. The successive treatment of the derivatized polymers with fluorescein isothiocyanate and a sulfhydryl-containing ligand (thioethanol, an MSH analog or a dynorphin analog) provided the desired composites. The polymer substrate was dialyzed extensively after each reaction step and the degree of substitution calculated spectrophotometrically. The peptide derivatives used for conjugation (Table 1) were synthesized by SPPS and purified by HPLC.

The binding of the fluorescent conjugates with a variety of cultured melanoma cells and other malignant and normal cell types as studied by fluorescence microscopy (Table 1) demonstrated that: a) the MSH-PVA conjugates bound

92-22135



599

2030310244

Best Available Copy

COMPONENT PART NOTICE

THIS PAPER IS A COMPONENT PART OF THE FOLLOWING COMPILATION REPORT:

TITLE: Peptides. Chemistry and Biology: Proceedings of the American Peptide Symposium
(12th) Held in Cambridge, Massachusetts on 16-21 June 1991.

TO ORDER THE COMPLETE COMPILATION REPORT, USE AD-A256 113.

THE COMPONENT PART IS PROVIDED HERE TO ALLOW USERS ACCESS TO INDIVIDUALLY
AUTHORED SECTIONS OF PROCEEDING, ANNALS, SYMPOSIA, ETC. HOWEVER, THE COMPONENT
SHOULD BE CONSIDERED WITHIN THE CONTEXT OF THE OVERALL COMPILATION REPORT AND
NOT AS A STAND-ALONE TECHNICAL REPORT.

THE FOLLOWING COMPONENT PART NUMBERS COMPRISE THE COMPILATION REPORT:

AD#: AD-P008 306 AD#: AD-P008 592
AD#: _____ AD#: _____
AD#: _____ AD#: _____

Accession For	
NTIS Card	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Table 1 Results demonstrating specific fluorescence labeling of melanotropin receptors on various melanoma cell types by fluorescent MSH-macromolecular composites

Conjugate ^a	Malignant cell types									Normal cell types	
	Mouse melanoma		Human							Mouse	
			Melanoma					Breast cancer			
	B-16	B-16 ^a	JH ^b	LR 1649	LR 1650	LR 1714 ^b	WC ^b	MA ^c	MCF-7	Spleen	Liver
FITC-PVA	-	-	-	-	-	-	-	-	-	-	-
FITC-PVA-S-MSH	+	+	+	+	+	+	+	+	-	-	-
FITC-PVA-S-S-MSH	+			+	+	+					
FITC-PVA-S-S-MSH + DTT	-			-	-	-					
FITC-PVA-S-MSH + DTT	+			+	+	+					
FITC-PVA-TE	-		-	-	-	-			-		
FITC-PVA-S-TE	-			-	-	-					
FITC-PVA-S-DYN	-	-									

(+) Indicates fluorescence labeling.

(-) Indicates no fluorescence labeling.

^a Cells prefixed in 1% formalin.

^b Amelanotic cell lines.

^c Similar positive responses were obtained with all other human melanoma cells that were assayed.

^d Abbreviations:

DTT = Dithiothreitol.

FITC = Fluorescein.

PVA = Poly vinyl alcohol.

S-DYN = Dynorphin analog; [Lys(β -thiopropionyl)³]Dynorphin(1-13)-NH₂.

S-MSH = MSH analog; N^o-des acetyl-N^o-thiopropionyl[Nle⁴, D-Phe⁷]- α -MSH.

TE = Thioethanol.

to all the melanoma cell lines but not to MCF-7 and to several normal cell types used as controls; b) conjugates lacking the MSH analog, or instead containing a dynorphin analog or thioethanol moieties did not bind melanoma cells; c) the treatment of FITC-PVA-S-S-MSH with DTT prior to its use in fluorescence labeling experiment abolished its ability to bind the melanoma cells. All these observations strongly supported the specificity of the binding between fluorescent MSH-PVA conjugates and melanoma cells.

Acknowledgements

This work was supported in part by a research grant from the U.S. Public Health Service (DK-17420 to VJH).

References

- Schwyzler, R. and Kriwaczek, V.M., Biopolymers, 20 (1981) 2011.
- Wunderlin, R., Sharma, S.D., Minakakis, P. and Schwyzler, R., Helv. Chim. Acta, 68 (1985) 12.